



















The Nobivac® recommended vaccination regime for cats

For cats that are not at risk of *Chlamydophila felis*, Intervet suggests the following regime:

8-9 weeks	 	Nobivac Tricat and Nobivac FeLV
12 weeks	 	Nobivac Tricat and Nobivac FeLV
Thereafter annually	 	Nobivac Tricat and Nobivac FeLV

For cats at risk of *Chlamydophila felis*, Intervet suggests the following regime:

8-9 weeks	 	Nobivac Forcat and Nobivac FeLV
12 weeks	 	Nobivac Forcat and Nobivac FeLV
Year 1	 	Nobivac Forcat and Nobivac FeLV
Year 2	 	Nobivac Ducat and Nobivac FeLV
Year 3	 	Nobivac Ducat and Nobivac FeLV
Year 4	 	Nobivac Tricat + Nobivac FeLV
etc, etc		

For cats living alone and strictly indoors:

8-9 weeks		Nobivac Tricat
12 weeks		Nobivac Tricat
Thereafter annually		Nobivac Tricat

If cats visit shows, go into boarding catteries, or are in contact with dogs, annual vaccination with **Nobivac Bb for Cats** should be added to the vaccination schedule.

Nobivac Tricat and Nobivac FeLV are fully licensed to be mixed. No information is available on the safety and efficacy from the concurrent use of Nobivac Forcat with any other. It is therefore recommended that no other vaccines should be administered within 14 days before or after vaccination with this product.